

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES


Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**



DOCKET NO. ORT-1230


IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

Applicant: S. Dax *et al.*
Serial No.: 09/552,969 Art Unit : 1624
Filed : April 20, 2000 Examiner: Hong Liu
Title : 3a,4,5,9b-tetrahydro-1h-benz[e]indol-2yl amine-derived neuropeptide y
receptor ligands useful in the treatment of obesity and other disorders

I hereby certify that this correspondence is being deposited with the
United States Postal Service as first class mail in an envelope addressed
to: Commissioner of Patents and Trademarks, Washington D.C. 20231 on

February 11, 2004
(Date of Deposit)

Ralph R. Palo
Name of Applicant, assignee, or Registered Representative


(Signature)

February 11, 2004
(Date of Signature)

Commissioner for Patents
P.O. Box 1450
Alexandria, VA 22313-1450

DECLARATION UNDER 37 C.F.R. 1.131

Dear Sir:

1. We, Scott L. Dax and James McNally declare that we are the inventors of the invention described and claimed in U.S. patent application Serial No. 09/522,969, filed on April 20, 2000, which application is based on provisional application Serial No. 60/132,660, filed May 5, 1999, and now abandoned.

2. We are presently, and were at and before completion of the invention, in the employ of Johnson & Johnson, which is the parent company of Ortho-McNeil Pharmaceutical Corporation, the assignee of record of the entire right, title and interest in the above-identified application.

3. We are familiar with the Office Action dated June 6, 2003 in which the above referenced application was rejected over a publication by McNally et al. which appeared in Bioorganic and Medicinal Chemistry Letters 10 (2000) 213-216.

4. We declare that the invention described and claimed in the above identified application was conceived by us in this country prior to February 7, 2000 and that such conception was coupled with due diligence by us in this country from just prior to February 7, 2000 to a reduction to practice of the invention.

5. Exhibits A-E attached hereto consist of true copies from which the dates have been removed of documents from the Ortho-McNeil Pharmaceutical Corporation research laboratories which record the generic conception of the invention.

6. Exhibits F-AA attached hereto consist of true copies from which the dates have been removed of documents from the Crystallitics Company which illustrate the reduction to practice of the invention.

7. Exhibits AB-AQ attached hereto consist of true copies from which the dates have been removed of documents from the Ortho-McNeil Pharmaceutical Corporation research laboratories which illustrate the operativeness of the invention. The assays and methods employed are known procedures which were published prior to February 7, 2000.

8. All of the above procedures and tests were carried out by us or at our direction from a period prior to February 7, 2000 to a reduction to practice of the invention. The results of the testing indicated that the compounds were useful in the treatment of obesity and other disorders.

9. We, Scott L. Dax and James McNally, further declare that all statements made herein of our own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Date: 2/10/2004

Date: 2/10/2004



SCOTT L. DAX



JAMES MCNALLY

Project No. 1227


Protocol / Experiment No.

Date

Subject

Purpose

Contd. from page



N.Wt 495.64

481.66 / 518.13

Wt 1.6g

1.38

mmol 3.22

16.1

Vol

16.1

from IPA/IPA/HCl

Lit in Et₂O

HPLC 3.306 ~ 75% 3.40 impurities 8.44

MS 482 MH⁺

Procedure

In a 200 mL RBF (mag stir, cond., Ar, heat mantle) 1.6g was added in portions to a solution of LAH (1.0M in THF 16.1 mL) in THF 20 mL at rt. Considerable foaming. Rxn heated to reflux 45 min., cooled in ice bath quenched with water 0.65 mL in THF ~ 5 mL, 10% NaOH 0.65 mL then Water 2.0 mL. SE in rt 1/2 h. Added Na₂SO₄ stir 1 h. Filter through celite - wash with THF then DCM. Combined organics stripped, residue taken up in IPA ~ 20 mL treated with HCl/IPA. No crystallization. ^{Solvent} Sample was evap under a stream of N₂. Residue triturated in Et₂O to give a pale pink solid 14794-65-1 1.38g

NMR, IR, MS, HPLC

Contd. on page 67

Investigator

Date

Read and Understood

Date

Project No. 11722




Procedure: Experiment 1b

Date _____

Subject:

Prooves

Contd. from page



	<u>1</u>	<u>2</u>	<u>3</u>	<u>4</u>	<u>5</u>
M.Wt	252.75	279.34	379.25	129.25	495.64
wt	1.0	1.16	1.58	1.79	1.87 _g
amp	3.96	4.15	4.15	13.8	4.22 3.7
Vol				2.41	
den				0.742	496 m.

HPLC 3.214

Agenda

In a 100 mL RBF (mag stir, Ar) DIER 2.4
was solvaged into a solution of 3.46 g &
3.65 g in DMF (5 mL) at rt. Fm stirred
for 15 min then cooled to 0°C added
1.1 g. Stirred 10°C for 1 hr. poured
products. washed with 5 mL of 1:1 EtOAc:Hex.
Rins stirred for 1 hr. then 1:1 Hex:EtOAc.
Hex evaporated. St. residue. St. dried to
4.6 g. 1.5 h. 100°C. 3.2 g. from 1.5 g. product.
at 100°C. 1.5 h. Rins. poured 1:1 Hex:EtOAc.
1.5 h. 100°C. 3.2 g. from 1.5 g. product.
washed in ether.
with 1:1 Hex:EtOAc to give 14794-62-1
1.67 g. MS✓ NPLC✓ NMR✓

Contd. on next:-

Investigator

Date _____

Learn, and Understood

Date _____

Project No. 1222

Protocol / Experiment No.

Date



Subject

Purpose

RWJ 355295-002-A

JMC

Contd. from page

1234

MW 215.25

77.08

62.84

216.29/252.75

Wt 3.5g

18.8

5.11

mmol 16.2

0.24 mol

81.3

Vol

50ml

HPLC 1.99 ~ 99%

Procedure

In a 200 ml RBF (mag stir, Ar) a solution of 1 3.5g & 2 18.8g in MeOH 50ml was stirred at 15 min. Added 3 5.11g & heated reflux 50 min. HPLC - rxn complete Rxn conc.

in vacuo at rt. - Treated with NaOH/H₂O

12g in 100 ml at 0°C. Pale grey solid collected

by filtration, wash H₂O Air dried - trit. in Et₂O

filter - air dry to give the free base 14794-61-FB

3.5g - removed 1/2 g store freezer

Remainder dissolved in THF/MeOH ~ 9/1 ~ 75 ml.

(need gentle heat to dissolve) add 1N HCl/Et₂O ~ 40 ml

cool on ice bath filter wash Et₂O air dry

14794-61-1 HCl salt 1.571 g

Contd. on page

Investigator

Date

Read and Understood

Date

Project No. 1227


Protocol / Experiment No.

Date

Subject

Purpose

Contd. from page 59



	<u>1</u>	2	3	4	5	6
MWt						215.25
wt						3.3g
amp						
Vol		60	50	5	100	HPLC
						2.864

Procedure

In a 500 ml RBF (mag stir, Ar) 1 (see p. 69) was taken up in DCM 60ml, H₂O 50ml, AcOH 5ml & MeOH 100ml. Resultant soln stirred rt 18h. Rxn complete by LC. Partitioned between DCM 100ml, H₂O 200ml. wash extr. Ar with DCM ~ 100ml. Combined organics wash w H₂O 2 x 100ml, sat'd NaHCO₃ 100ml - dried over Mg 60y & stripped to give a brown oil 14794-60-3.3, HPLC ✓

Contd. on page

Investigator

Date

Read and Understood

Date

Project No. 1227


Protocol / Experiment No.

Date

Subject

Purpose see 14139-173 Ser NMR 3

Contd. from page



Procedure

In a 300ml RBF (Ar, mag stir) 2 3.5ml was added to a solution of 2 5.64g in MeOH 60ml. Rxn stirred rt 1.5h. Solid ppt'd. Rxn mix cool in IPA/ice bath. Solid collected by filtration, wash cold MeOH and dried. 14794-59-1 HPLC ✓

Solid taken up in CH₃CN ~ 60ml (mag stir, Ar) treated with 4 2.21ml stir rt 1h. Solid ppt collected by filtration - wash with CH₃CN, Et₂O - 14794-59-2 HPLC

0.990 2 2.812 M+ 269

Second crop (HPLC ✓) collected & used with 14794-59-2 on p. 60

Contd. on page

60

Investigator

Date


Read and Understood

Date

Jim McNally

Exhibit F

CRYSTAL STRUCTURE ANALYSIS REPORT and TABLES for




Crystalytics Company Reference Code: CFS4-0799

For: The R. W. Johnson Research Institute

Sample #14794-65-1

Jim McNally

Copy 2



CRYSTALYTICS COMPANY

CRYSTAL STRUCTURE CONSULTING

P.O. BOX 82286 • LINCOLN, NEBRASKA 68501
TELEPHONE (402) 421-2797


**BRIEF EXPERIMENTAL DESCRIPTION TO BE INCLUDED IN TEXT OR
AS A FOOTNOTE AT TIME OF PUBLICATION**

Single crystals of $[C_{27}H_{36}N_3O_3S][Cl]$ are, at $20 \pm 1^\circ C$, orthorhombic, space group $Pna2_1 - C_{2v}^9$ (No. 33) with $a = 42.245(1) \text{ \AA}$, $b = 5.3262(2) \text{ \AA}$, $c = 11.9748(4) \text{ \AA}$, $V = 2694.4(2) \text{ \AA}^3$, and $Z = 4$ $\{d_{\text{calcd}} = 1.277 \text{ gcm}^{-3}$; $\mu_a(\text{CuK}\alpha) = 2.24 \text{ mm}^{-1}\}$. A full hemisphere of diffracted intensities (omega or phi scans with width of 0.25°) was measured using graphite-monochromated $\text{CuK}\alpha$ radiation on a Siemens X-1000 HI-STAR Multiwire area detector. X-rays were provided by a Siemens M18XHF rotating anode operated at 40kV and 70mA. The sample was a nonmerohedrally twinned specimen containing two domains; the major domain had approximately 17 times the volume of the minor domain and only 4% of the reflections for the two domains were partially overlapped. There were no totally overlapping reflections and partially overlapping reflections were not used for structure refinement. The diffraction data from each domain was used to solve the structure independently and both gave the same species. The results reported herein are for the major domain but nearly identical structural parameters resulted from refinement using the nonoverlapping data for both domains simultaneously. Structure refinement using just the data from the minor domain gave structural parameters with reduced precision.

Lattice constants for the major domain were determined with the Siemens SAINT software package using peak centers for 2292 reflections. A total of 4902 integrated reflection intensities having $2\theta(\text{CuK}\alpha) < 104.3^\circ$ were produced using the Siemens program SAINT. A total of 2788 of these were independent and gave $R_{\text{int}} = 0.057$. The Siemens SHELXTL-PC software package was used to solve the structure using "Direct Methods" techniques. All stages of weighted full-matrix least-squares refinement were conducted using F_o^2 data and the SHELXTL-PC Version 5 software package and converged to give R_1 (unweighted, based on F) = 0.053 for 2223 independent absorption-corrected reflections having $2\theta(\text{CuK}\alpha) < 104.3^\circ$ and $I > 2\sigma(I)$ and wR_2 (weighted, based on F^2) = 0.145 for 2673 independent absorption-corrected reflections having $2\theta(\text{CuK}\alpha) < 104.3^\circ$ and $I > 0$. Final R values for all 2788 independent absorption-corrected reflections having $2\theta(\text{CuK}\alpha) < 104.3^\circ$ are: R_1 (unweighted, based on F) = 0.074 and wR_2 (weighted, based on F^2) = 0.166. The structural model incorporated anisotropic thermal parameters for all nonhydrogen

Exhibit H

atoms and isotropic thermal parameters for all hydrogen atoms. Hydrogen atoms H_{1N}, H_{2N} and H_{3N} were located from a difference Fourier map and refined as independent isotropic atoms. The remaining hydrogen atoms were included in the structure factor calculations as idealized atoms (assuming sp²- or sp³-hybridization of the carbon atoms and C-H bond lengths of 0.93-0.98 Å) "riding" on their respective carbon atoms. The isotropic thermal parameters for H_{1N}, H_{2N} and H_{3N} refined to final U_{iso} values of 0.10(3), 0.11(3) and 0.01(2) Å², respectively. The isotropic thermal parameters of the remaining hydrogen atoms were fixed at values 1.2 (nonmethyl) or 1.5 (methyl) times the equivalent isotropic thermal parameters of the carbon atoms to which they are covalently bonded. The methyl group (C₂₇ and its hydrogens) was refined as a rigid rotor (using idealized sp³-hybridized geometry and a C-H bond length of 0.96 Å) with three rotational parameters in each least-squares cycle. The refined values of these rotational parameters gave O-C-H angles which ranged from 105° to 117°.



Crystalytics Company Crystal Structure Analysis Report

Compound Formula: $[C_{27}H_{36}N_3O_3S][Cl]$

Reference Code: CFS4-0799

R. W. J. sample #14794-65-1 (Jim McNally)

Description of Single-Crystal Sample and Mounting Used for Data Collection:

- 1) Color: Colorless
- 2) Shape: Flat plate
- 3) Dimensions: 0.025 mm. x 0.125 mm. x 0.175 mm.
- 4) Indices of Faces:
- 5) Crystal Mount: Crystal was glued with epoxy to the end of a thin glass fiber.
- 6) Crystal Orientation: Crystal was oriented with its longest edge nearly parallel to the ϕ axis of the diffractometer.
- 7) Comments: Sample was recrystallized from methanol/ethylacetate solution.

Space Group and Cell Data:

- 1) Crystal System: Orthorhombic Space Group and Number¹: $Pna2_1 - C_{2v}^9$ (No. 33)
- 2) Number of Computer-Centered Reflections Used in the Least-Squares Refinement of the Cell Dimensions: 2292 measured at: $20 \pm 2^\circ C$
- 3) Lattice Constants with esd's:

$a = 42.245(1) \text{ \AA}$	$\alpha = 90.000^\circ$	$V = 2694.4(2) \text{ \AA}^3$
$b = 5.3262(2) \text{ \AA}$	$\beta = 90.000^\circ$	$Z = 4 \text{ formula units}$
$c = 11.9748(4) \text{ \AA}$	$\gamma = 90.000^\circ$	$\lambda = 1.54178 \text{ \AA}$
- 4) Molecular Weight: 518.10 amu/formula unit Calculated Density: 1.277 g cm^{-3}
- 5) Linear Absorption Coefficient^{2a}: 2.24 mm^{-1} $F(000) = 1104.$
- 6) Comments: The sample was a nonmerohedrally twinned specimen containing two domains; the major domain had approximately 17 times the volume of the minor domain and only 4% of the reflections for the two domains were partially overlapped. There were no totally overlapping reflections and partially overlapping reflections were not used for structure refinement. The diffraction data from each domain was used to solve the structure independently and both gave the same species. The results reported herein are for the major domain but nearly identical structural parameters resulted from refinement using the nonoverlapping data for both domains simultaneously. Structure refinement using just the data from the minor domain gave structural parameters with reduced precision.

Description of Data Collection³:

- 1) Instrument: Bruker X-1000 HI-STAR Single Crystal Multiwire Diffraction System
- 2) X-ray Source: Bruker M18XHF Rotating Anode with 0.3 x 3.0 mm. filament
- 3) Radiation: CuK α Power: 40 kV 70 mA
- 4) X Monochromator: X Graphite Other (Specify:)
 Filter: Nickel Niobium Other (Specify:)
- 5) Incident Beam Collimator Diameter: 0.5 mm Temperature: 20 \pm 2 $^{\circ}$ C
- 6) Scan Axis: X Omega or X Phi
- 7) Scan Width: 0.25 $^{\circ}$ 2 θ Range of Data : 8.38 $^{\circ}$ - 104.28 $^{\circ}$
- 8) Sample to Detector Distance: 8.56 cm
- 9) Portion of Ewald Sphere Collected: Hemisphere
- 10) Number of frames collected: 3228 Seconds/frame: 60
- 11) Total Number of Reflections Collected: 4902
- 12) Number of Independent Reflections Collected: 2788
- 13) Data Collected: $-28 \leq h \leq 43$; $-5 \leq k \leq 5$; $-11 \leq l \leq 12$ $R_{\text{int}}^4 = 0.057$

Data Reduction³:

- 1) Lorentz and Polarization Corrections? Yes
- 2) Absorption Correction: Yes Range of transmission factors: 0.524 - 0.666
XX Empirical Correction using Measurements for Equivalent Reflections
(508 Reflections used)
 Face-Indexed Gaussian Grid Correction
- 3) Comments:

Structure Solution⁵:

- 1) Method(s) Used in Structure Solution
 Heavy-atom Patterson Techniques
XX Direct Methods
a) XX SHELXTL/PC
b) Other
 Other Techniques
- 2) Hydrogen Atom Positions Located? Yes
After Refinement Cycle # 2 by XX Difference Fourier
XX Calculated
- 3) Comments:

Structure Refinement⁵: (see next page for summary of refinement cycles)

1) Final Scale Factor: 0.327(1)

2) Extinction Parameter⁶ Refined? Yes Final Value: 0.0002(2)Form: $k[1 + 0.001(x)(F_c^2)(\lambda^3)/\sin(2\theta)]^{-1/4}$ 3) Anomalous Dispersion Corrections^{2b} for Which Atoms: Cl, S, O, N, C

4) Variable Occupancies for Which Atoms? None

Atom	Final Occupancy	Atomic Form Factor ^{2c} Used
------	-----------------	---------------------------------------

5) Refinement Constraints/Restraints: Hydrogen atoms H_{1N}, H_{2N} and H_{3N} were located from a difference Fourier map and refined as independent isotropic atoms. The remaining hydrogen atoms were included in the structure factor calculations as idealized atoms (assuming sp²- or sp³- hybridization of the carbon atoms and C-H bond lengths of 0.93-0.98 Å) "riding" on their respective carbon atoms. The isotropic thermal parameters for H_{1N}, H_{2N} and H_{3N} refined to final U_{iso} values of 0.10(3), 0.11(3) and 0.01(2) Å², respectively. The isotropic thermal parameters of the remaining hydrogen atoms were fixed at values 1.2 (nonmethyl) or 1.5 (methyl) times the equivalent isotropic thermal parameters of the carbon atoms to which they are covalently bonded. The methyl group (C₂₇ and its hydrogens) was refined as a rigid rotor (using idealized sp³- hybridized geometry and a C-H bond length of 0.96 Å) with three rotational parameters in each least-squares cycle. The refined values of these rotational parameters gave O-C-H angles which ranged from 105° to 117°.

6) Shift/Error Analysis for Final Least-Squares Cycle⁷:Maximum Shift for all Parameters: 0.000 σ_p Mean Shift for all Parameters: 0.000 σ_p

7) Peaks found in Final Difference Fourier Map: There were no peaks present in the final difference Fourier map above the background level (0.35 e⁻/Å³). The minimum and mean electron density in the final difference Fourier were -0.19 and 0.00 e⁻/Å³, respectively. The rms deviation from the mean electron density was 0.04 e⁻/Å³.

Summary of Full Matrix Least-Squares Refinement⁸ Cycles

Cycle Number	sin θ / λ		Anisotropic ⁹ Atoms Number and Type	Isotropic Atoms			R_1 (unweighted, based on F)		$F_o / \sigma(F_o)$ Cutoff	R_2^{12}	Total # Independent Reflections	'Goodness- of-fit' (Goof) ¹³	Extincti n Correcti n
	Minimum	Maximum		Number and Type	Positions Refined	Thermal Parameters	# Refined Parameters	R_1^{11}					
1	0.00	0.51		27 C, 3 N 3 O, 1 S 1 Cl	X	X	141	0.093	4.0	0.249	2673	1.755	
2	0.00	0.51	27 C, 3 N 3 O, 1 S 1 Cl				316	0.074	4.0	0.193	2673	1.400	
3	0.00	0.51	27 C, 3 N 3 O, 1 S 1 Cl	*36H	X	X	332	0.053	4.0	0.145	2673	1.045	X

* See Item 5 on page 3 regarding the treatment of the hydrogen atoms.

Final Statistics from Cycle #3 for All of the Reflection Data: $R_1 = 0.074$; $wR_2 = 0.166$; GOOF = 1.169 for 2788 reflections

The correctness of the assigned absolute configuration was checked using the "Flack" absolute structure parameter¹⁴ which refined to a final value of 0.18(3).

References and Notes

1. "International Tables for X-Ray Crystallography", Vol. A, Kluwer Academic Publishers, Dordrecht, 1995.
2. "International Tables for X-Ray Crystallography", Vol. C, Kluwer Academic Publishers, Dordrecht, 1992; a) Tables 4.2.4.2 pp. 193-199; b) Tables 4.2.6.8 pp 219-222; c) Tables 6.1.1.4 pp 500-502.
3. Data acquisition and reduction was accomplished using standard versions of Siemens/Bruker software for the diffraction system.
4. $R_{\text{int}} = \Sigma |F_o^2 - F_o^2(\text{mean})| / \Sigma [F_o]^2$
5. All structure determination and refinement calculations were performed on an IBM compatible 486 or 586 personal computer using the Siemens/Bruker SHELXTL Version 5.0 PC interactive software package.
6. A. C. Larson in "Crystallographic Computing", 1970, Ed. F. R. Ahmed, Munksgaard, Copenhagen, pp 291-294.
7. σ_p is the estimated standard deviation of the parameter in question.
8. Refinement on F^2 for all reflections except for 115 with negative F^2 . Weighted R-factors wR_2 and all goodnesses of fit S are based on F^2 , conventional R-factors R_1 are based on F , with F set to zero for negative F^2 . The observed criterion of $F^2 > 2\text{sigma}(F^2)$ is used only for calculating "R-factor obs" etc. and is not relevant to the choice of reflections for refinement. R-factors based on F^2 are statistically about twice as large as those based on F , and R-factors based on all data will be even larger.
9. The anisotropic thermal parameter is of the form:
 $\exp[-2\pi^2(U_{11}h^2a^{*2} + U_{22}k^2b^{*2} + U_{33}l^2c^{*2} + 2U_{12}hka^*b^* + 2U_{13}hla^*c^* + 2U_{23}klb^*c^*)]$.
10. The weighting scheme used is defined as: $w = 1 / [\sigma^2(F_o^2) + (a^*P)^2 + b^*P + d + e^*\sin(\theta)]$ where $P = [F_o^2 + 2F_c^2]/3$. In this case, $a = 0.0938$, $b = 0$, $d = 0$ and $e = 0$.
11. $R_1 = \Sigma ||F_o| - |F_c|| / \Sigma |F_o|$
12. $wR_2 = [\Sigma [w(F_o^2 - F_c^2)^2] / \Sigma [w(F_o^2)^2]]^{1/2}$
13. $\text{GooF} = S = [\Sigma [w(F_o^2 - F_c^2)^2] / (n-p)]^{1/2}$ where n is the total number of reflections and p is the number of parameters refined.
14. The value of the "Flack absolute structure parameter", x , should be 0.00 for the correct enantiomorphic description and 1.00 for the inverted description: a) H. D. Flack, *Acta Cryst.*, 1983, *A39*, 876-881; b) G. Bernardinelli and H. D. Flack, *Acta Cryst.*, 1985, *A41*, 500-511.

Exhibit N

Table 1. Atomic Coordinates for Nonhydrogen Atoms in Crystalline
 $[\text{C}_{27}\text{H}_{36}\text{N}_3\text{O}_3\text{S}][\text{Cl}]^a$

Atom	Fractional Coordinates			Equivalent Isotropic
Type ^b	10 ⁴ x	10 ⁴ y	10 ⁴ z	Thermal Parameter, U, Å ² x 10 ³ ^c
Cation				
S	1890(1)	-1817(3)	3378(2)	62(1)
O ₁	1739(1)	-4228(8)	3392(5)	84(1)
O ₂	2083(1)	-1058(9)	4292(4)	79(1)
O ₃	-1850(1)	7660(12)	3106(5)	105(2)
N ₁	-362(1)	2030(13)	2990(6)	62(2)
N ₂	-53(1)	5255(12)	2246(6)	66(2)
N ₃	1619(2)	209(16)	3278(7)	67(2)
C ₁	-297(2)	3645(14)	2220(6)	59(2)
C ₂	-525(2)	3482(13)	1278(5)	65(2)
C ₃	-793(2)	1762(14)	1736(6)	67(2)
C ₄	-1083(2)	3353(14)	2115(6)	64(2)
C ₅	-1248(2)	4655(17)	1347(7)	86(2)
C ₆	-1510(2)	6149(16)	1651(8)	87(3)
C ₇	-1598(2)	6274(16)	2739(8)	77(2)
C ₈	-1441(2)	4911(14)	3526(7)	73(2)
C ₉	-1175(1)	3462(11)	3245(6)	55(2)
C ₁₀	-996(2)	2072(16)	4119(6)	77(2)
C ₁₁	-829(2)	-256(15)	3679(6)	75(2)
C ₁₂	-629(2)	351(13)	2681(6)	61(2)
C ₁₃	186(2)	5209(14)	3088(6)	66(2)
C ₁₄	443(1)	3309(12)	2830(5)	53(2)
C ₁₅	630(2)	4008(13)	1781(6)	62(2)
C ₁₆	906(1)	2211(14)	1576(5)	60(2)
C ₁₇	1121(1)	2009(12)	2563(5)	52(2)
C ₁₈	931(1)	1224(13)	3608(5)	59(2)
C ₁₉	663(2)	3025(14)	3812(5)	64(2)

Table 1. (continued)

Atom Type ^b	Fractional Coordinates			Equivalent Isotropic Thermal Parameter, U, Å ² x 10 ³ ^c
	10 ⁴ x	10 ⁴ y	10 ⁴ z	
C ₂₀	1388(1)	186(14)	2353(6)	63(2)
C ₂₁	2110(2)	-1655(13)	2128(6)	60(2)
C ₂₂	2064(2)	-3338(17)	1289(8)	94(3)
C ₂₃	2237(2)	-3108(20)	305(9)	111(3)
C ₂₄	2456(2)	-1242(21)	189(9)	105(3)
C ₂₅	2502(2)	386(19)	1015(9)	102(3)
C ₂₆	2332(2)	215(17)	1990(8)	90(3)
C ₂₇	-1972(2)	9523(21)	2397(10)	122(4)
Anion				
Cl	-64(1)	744(4)	5362(2)	75(1)

^a The numbers in parentheses are the estimated standard deviations in the last significant digit.

^b Atoms are labeled in agreement with Figure 1.

^c This is one-third of the trace of the orthogonalized U_{ij} tensor.

Exhibit P

Table 2. Anisotropic Thermal Parameters for Nonhydrogen Atoms in Crystalline
 $[C_{27}H_{36}N_3O_3S][Cl]$ ^{a,b}

Atom Type ^c	Anisotropic Thermal Parameters ($\text{\AA}^2 \times 10^3$)					
	U_{11}	U_{22}	U_{33}	U_{23}	U_{13}	U_{12}
Cation						
S	59(1)	55(1)	74(1)	6(1)	-14(1)	-3(1)
O ₁	82(3)	48(3)	121(4)	21(3)	-10(3)	-13(2)
O ₂	81(3)	79(3)	77(3)	6(3)	-22(3)	-15(3)
O ₃	79(3)	116(5)	120(6)	20(4)	21(4)	29(3)
N ₁	58(4)	73(4)	56(4)	4(4)	0(3)	6(3)
N ₂	56(4)	56(4)	86(5)	4(4)	8(3)	4(3)
N ₃	68(4)	61(5)	73(5)	-11(5)	-12(3)	-17(4)
C ₁	64(5)	69(5)	44(4)	13(4)	1(4)	22(4)
C ₂	62(4)	74(5)	60(4)	4(4)	8(4)	16(4)
C ₃	51(4)	71(5)	79(5)	-15(4)	15(4)	12(4)
C ₄	61(4)	66(5)	66(5)	8(4)	-11(4)	-10(4)
C ₅	78(5)	108(7)	74(5)	3(5)	-15(5)	18(5)
C ₆	68(5)	96(7)	96(7)	27(5)	-10(5)	19(5)
C ₇	41(4)	95(6)	95(6)	19(5)	0(4)	10(4)
C ₈	60(4)	76(5)	84(5)	4(5)	7(4)	-8(4)
C ₉	47(3)	57(4)	60(5)	9(4)	1(3)	1(3)
C ₁₀	69(5)	91(6)	72(5)	10(5)	7(4)	-12(5)
C ₁₁	67(4)	80(5)	77(5)	10(5)	-15(4)	-6(4)
C ₁₂	56(4)	56(4)	72(5)	-5(4)	-3(4)	4(4)
C ₁₃	55(4)	68(4)	75(5)	-6(4)	3(4)	2(3)
C ₁₄	47(3)	52(4)	60(4)	2(3)	7(3)	7(3)
C ₁₅	57(4)	60(4)	69(5)	2(4)	4(4)	3(4)
C ₁₆	54(4)	67(5)	58(4)	-1(4)	9(3)	7(3)
C ₁₇	50(4)	54(4)	52(4)	0(3)	4(3)	0(3)
C ₁₈	52(4)	72(5)	53(4)	0(4)	0(3)	3(3)
C ₁₉	66(4)	80(5)	47(4)	3(4)	6(3)	-13(4)
C ₂₀	50(4)	69(5)	68(5)	-9(4)	-2(4)	7(4)

Exhibit Q

Table 2. (continued)

Atom Type ^c	Anisotropic Thermal Parameters ($\text{\AA}^2 \times 10^3$)					
	U_{11}	U_{22}	U_{33}	U_{23}	U_{13}	U_{12}
C ₂₁	57(4)	47(4)	75(5)	3(4)	-1(4)	0(4)
C ₂₂	84(6)	87(6)	109(7)	-34(6)	11(5)	-8(5)
C ₂₃	105(7)	110(8)	118(8)	-33(7)	40(7)	0(6)
C ₂₄	104(7)	104(7)	108(8)	-21(7)	27(6)	7(6)
C ₂₅	96(6)	100(7)	109(8)	-7(7)	33(6)	-23(6)
C ₂₆	92(6)	80(6)	99(7)	-12(5)	28(5)	-21(5)
C ₂₇	97(7)	103(8)	164(11)	14(8)	-6(7)	33(6)
			Anion			
Cl	85(1)	75(1)	65(1)	11(1)	-10(1)	-1(1)

- ^a The numbers in parentheses are the estimated standard deviations in the last significant digit.
- ^b The form of the anisotropic thermal parameter is: $\exp[-2\pi^2(U_{11}h^2a^{*2} + U_{22}k^2b^{*2} + U_{33}l^2c^{*2} + 2U_{12}hka^*b^* + 2U_{13}hla^*c^* + 2U_{23}klb^*c^*)]$.
- ^c Atoms are labeled in agreement with Figure 1.

Exhibit R

Table 3. Atomic Coordinates for Hydrogen Atoms in Crystalline
 $[C_{27}H_{36}N_3O_3S][Cl]^a$

Atom Type ^b	Fractional Coordinates		
	10^4x	10^4y	10^4z
H _{1N} ^c	-288(20)	1722(159)	3622(80)
H _{2N} ^c	1(18)	6723(161)	1614(80)
H _{3N} ^c	1657(12)	1194(90)	3415(53)
H _{2b}	-427	2747	623
H _{2c}	-607	5129	1086
H ₃	-858	573	1155
H ₅	-1187	4570	601
H ₆	-1622	7037	1110
H ₈	-1511	4938	4262
H _{10a}	-1141	1579	4707
H _{10b}	-839	3188	4445
H _{11a}	-696	-957	4263
H _{11b}	-985	-1509	3476
H ₁₂	-542	-1217	2382
H _{13a}	89	4797	3800
H _{13b}	279	6865	3152
H ₁₄	342	1683	2697
H _{15a}	490	3978	1140
H _{15b}	711	5703	1859
H _{16a}	824	561	1397
H _{16b}	1027	2795	938
H ₁₇	1213	3668	2704
H _{18a}	847	-455	3504
H _{18b}	1070	1200	4253
H _{19a}	750	4658	3997
H _{19b}	542	2449	4450
H _{20a}	1302	-1493	2271
H _{20b}	1494	628	1662

Table 3. (continued)

Atom Type ^b	Fractional Coordinates		
	10 ⁴ x	10 ⁴ y	10 ⁴ z
H ₂₂	1918	-4630	1371
H ₂₃	2204	-4234	-276
H ₂₄	2572	-1109	-467
H ₂₅	2652	1655	931
H ₂₆	2367	1368	2561
H _{27a}	-2103	10560	2860
H _{27b}	-1822	10551	2008
H _{27c}	-2101	8649	1868

- ^a Hydrogen atoms covalently bonded to nitrogens (H_{1N}, H_{2N} and H_{3N}) were located from a difference Fourier map and refined as independent isotropic atoms. The remaining hydrogen atoms were included in the structure factor calculations as idealized atoms (assuming sp²- or sp³-hybridization of the carbon atoms and C-H bond lengths of 0.93 Å to 0.98 Å) "riding" on their respective carbon atoms. The methyl group (C₂₇ and its hydrogens) was refined as a rigid rotor (using idealized sp³-hybridized geometry and a C-H bond length of 0.96 Å) with three rotational parameters in each least-squares cycle. The refined values of these rotational parameters gave O-C-H angles which ranged from 105° to 117°. The isotropic thermal parameters for H_{1N}, H_{2N} and H_{3N} refined to final U_{iso} values of 0.10(3), 0.11(3) and 0.01(2) Å², respectively. The isotropic thermal parameters of the remaining hydrogen atoms were fixed at values 1.2 (nonmethyl) or 1.5 (methyl) times the equivalent isotropic thermal parameters of the carbon atoms to which they are covalently bonded.
- ^b Hydrogen atoms which are covalently bonded to carbon are labeled with the same numerical subscript(s) as their carbon atoms with an additional literal subscript (a, b or c) where necessary to distinguish between hydrogens bonded to the same carbon atom. The amine hydrogen atoms are labeled H_{1N} and H_{2N} and the sulfonamide hydrogen atom is labeled H_{3N}.
- ^c The numbers in parentheses are the estimated standard deviations in the last significant digit.

Exhibit T

Table 4. Bond Lengths in Crystalline $[\text{C}_{27}\text{H}_{36}\text{N}_3\text{O}_3\text{S}][\text{Cl}]$ ^a

Type ^b	Length, Å	Type ^b	Length, Å
S-O ₁	1.434(4)	S-O ₂	1.423(5)
S-N ₃	1.579(8)	S-C ₂₁	1.763(7)
O ₃ -C ₇	1.368(9)	O ₃ -C ₂₇	1.404(11)
N ₁ -C ₁	1.291(9)	C ₁ -C ₂	1.487(9)
N ₂ -C ₁	1.340(9)	C ₉ -C ₁₀	1.489(10)
N ₁ -C ₁₂	1.487(9)	N ₁ -H _{1N}	0.84(9)
N ₂ -C ₁₃	1.425(9)	N ₂ -H _{2N}	1.11(9)
N ₃ -C ₂₀	1.477(10)	N ₃ -H _{3N}	0.57(5)
C ₂ -C ₃	1.555(9)	C ₄ -C ₅	1.346(10)
C ₃ -C ₁₂	1.525(10)	C ₄ -C ₉	1.408(10)
C ₃ -C ₄	1.557(10)	C ₅ -C ₆	1.413(11)
C ₁₀ -C ₁₁	1.520(10)	C ₆ -C ₇	1.356(12)
C ₁₁ -C ₁₂	1.498(10)	C ₇ -C ₈	1.362(10)
C ₁₃ -C ₁₄	1.517(9)	C ₈ -C ₉	1.406(9)
C ₁₄ -C ₁₉	1.505(9)	C ₂₁ -C ₂₂	1.361(10)
C ₁₄ -C ₁₅	1.531(9)	C ₂₁ -C ₂₆	1.377(10)
C ₁₅ -C ₁₆	1.529(9)	C ₂₂ -C ₂₃	1.392(13)
C ₁₆ -C ₁₇	1.493(9)	C ₂₃ -C ₂₄	1.364(13)
C ₁₇ -C ₂₀	1.509(9)	C ₂₄ -C ₂₅	1.329(12)
C ₁₇ -C ₁₈	1.544(9)	C ₂₅ -C ₂₆	1.375(12)
C ₁₈ -C ₁₉	1.504(9)		

^a The numbers in parentheses are the estimated standard deviations in the last significant digit.

Exhibit U

b Atoms are labeled in agreement with Figure 1.

Exhibit V

Table 5. Bond Angles in Crystalline $[\text{C}_{27}\text{H}_{36}\text{N}_3\text{O}_3\text{S}][\text{Cl}]^a$

Type ^b	Angle, (deg)	Type ^b	Angle, (deg)
O_2SO_1	120.0(3)	O_2SC_{21}	109.8(3)
O_2SN_3	106.2(4)	O_1SC_{21}	106.8(4)
O_1SN_3	106.8(3)	N_3SC_{21}	106.5(4)
$\text{C}_7\text{O}_3\text{C}_{27}$	118.2(7)	$\text{C}_1\text{N}_2\text{C}_{13}$	123.2(7)
$\text{C}_1\text{N}_1\text{C}_{12}$	112.7(7)	$\text{C}_{20}\text{N}_3\text{S}$	122.1(6)
$\text{C}_1\text{N}_1\text{H}_{1\text{N}}$	134(6)	$\text{C}_{13}\text{N}_2\text{H}_{2\text{N}}$	111(4)
$\text{C}_{12}\text{N}_1\text{H}_{1\text{N}}$	113(6)	$\text{C}_{20}\text{N}_3\text{H}_{3\text{N}}$	114(6)
$\text{C}_1\text{N}_2\text{H}_{2\text{N}}$	126(4)	$\text{SN}_3\text{H}_{3\text{N}}$	114(6)
$\text{N}_1\text{C}_1\text{N}_2$	125.0(7)	$\text{N}_1\text{C}_1\text{C}_2$	111.4(7)
$\text{N}_2\text{C}_1\text{C}_2$	123.6(6)		
$\text{C}_1\text{C}_2\text{C}_3$	103.8(6)	$\text{C}_{19}\text{C}_{14}\text{C}_{15}$	110.3(5)
$\text{C}_{12}\text{C}_3\text{C}_2$	102.9(5)	$\text{C}_{13}\text{C}_{14}\text{C}_{15}$	112.0(5)
$\text{C}_{12}\text{C}_3\text{C}_4$	114.1(6)	$\text{C}_{16}\text{C}_{15}\text{C}_{14}$	112.0(5)
$\text{C}_2\text{C}_3\text{C}_4$	110.7(6)	$\text{C}_{17}\text{C}_{16}\text{C}_{15}$	112.4(5)
$\text{C}_9\text{C}_{10}\text{C}_{11}$	113.4(6)	$\text{C}_{16}\text{C}_{17}\text{C}_{20}$	111.6(5)
$\text{C}_{12}\text{C}_{11}\text{C}_{10}$	111.2(6)	$\text{C}_{16}\text{C}_{17}\text{C}_{18}$	110.3(5)
$\text{N}_1\text{C}_{12}\text{C}_{11}$	111.1(6)	$\text{C}_{20}\text{C}_{17}\text{C}_{18}$	110.4(5)
$\text{N}_1\text{C}_{12}\text{C}_3$	103.4(6)	$\text{C}_{19}\text{C}_{18}\text{C}_{17}$	110.5(5)
$\text{C}_{11}\text{C}_{12}\text{C}_3$	116.3(5)	$\text{C}_{18}\text{C}_{19}\text{C}_{14}$	113.7(5)
$\text{N}_2\text{C}_{13}\text{C}_{14}$	112.0(6)	$\text{N}_3\text{C}_{20}\text{C}_{17}$	111.4(6)
$\text{C}_{19}\text{C}_{14}\text{C}_{13}$	110.5(5)		
$\text{C}_5\text{C}_4\text{C}_9$	119.5(7)	$\text{C}_8\text{C}_9\text{C}_{10}$	120.8(7)

Exhibit W

Table 5. (continued)

Type ^b	Angle, (deg)	Type ^b	Angle, (deg)
C ₅ C ₄ C ₃	119.2(7)	C ₄ C ₉ C ₁₀	121.0(6)
C ₉ C ₄ C ₃	121.3(6)	C ₂₂ C ₂₁ C ₂₆	119.0(7)
C ₄ C ₅ C ₆	121.4(8)	C ₂₂ C ₂₁ S	121.3(6)
C ₇ C ₆ C ₅	119.3(7)	C ₂₆ C ₂₁ S	119.7(6)
C ₆ C ₇ C ₈	120.4(7)	C ₂₁ C ₂₂ C ₂₃	119.5(8)
C ₆ C ₇ O ₃	123.2(8)	C ₂₄ C ₂₃ C ₂₂	120.4(10)
C ₈ C ₇ O ₃	116.3(8)	C ₂₅ C ₂₄ C ₂₃	120.0(9)
C ₇ C ₈ C ₉	121.1(8)	C ₂₄ C ₂₅ C ₂₆	120.7(9)
C ₈ C ₉ C ₄	118.3(6)	C ₂₅ C ₂₆ C ₂₁	120.4(9)

^a The numbers in parentheses are the estimated standard deviations in the last significant digit.

^b Atoms are labeled in agreement with Figure 1.

Exhibit X

Table 6. Hydrogen-Bonding Interactions in Crystalline $[C_{27}H_{36}N_3O_3S][Cl]$

Donor Atom (D) ^a	Acceptor Atom (A)	Distance Å D...A	Distance Å H...A	Angle deg. D-H...A	Angle deg. H-D...A	Angle deg. H...A-X ^b	Asymmetric Unit of A ^c
N ₁ -H _{1N}	Cl	3.183	2.35	178	2	124(H _{2N})	x, y, z
N ₂ -H _{2N}	Cl	3.142	2.03	174	4	124(H _{1N})	-x, 1-y, -0.5+z
N ₃ -H _{3N}	O ₁	3.009	2.46	161	16	162(S)	x, 1+y, z

^a The hydrogen atom involved in the interaction is also indicated.

^b The symbol X is used to denote the atoms which are covalently or hydrogen bonded to the acceptor atoms.

^c All donor atoms belong to the asymmetric unit for which fractional atomic coordinates are given in Tables 1 and 3.

FIGURE CAPTIONS

- Figure 1a. shows a perspective drawing of the solid-state structure for the HCl salt of $[C_{27}H_{35}N_3O_3S]$. Nonhydrogen atoms are represented by 50% probability thermal vibration ellipsoids and hydrogen atoms are represented by arbitrarily-small spheres which are in no way representative of their true thermal motion. The hydrogen-bonding interaction between the Cl^- anion and the hydrogen on amine nitrogen N_1 is shown with a dashed line.
- Figure 1b. shows a perspective drawing of the solid-state structure for the HCl salt of $[C_{27}H_{35}N_3O_3S]$. The view is the same as in Figure 1a but the chlorine atom is now represented as a large dotted sphere, oxygen and nitrogen atoms are now represented by medium-sized shaded spheres, and sulfur, carbon and hydrogen atoms are represented by large, medium and small open spheres, respectively. The hydrogen-bonding interaction between the Cl^- anion and the hydrogen on amine nitrogen N_1 is shown with a dashed line.
- Figure 1c. shows a perspective drawing of the structure for the $[C_{27}H_{36}N_3O_3S]^+$ cation, as observed in the solid-state structure of its Cl^- salt. Nonhydrogen atoms are represented by 50% probability thermal vibration ellipsoids and hydrogen atoms are represented by arbitrarily-small spheres which are in no way representative of their true thermal motion.
- Figure 1d. shows a perspective drawing of the structure for the $[C_{27}H_{36}N_3O_3S]^+$ cation, as observed in the solid-state structure of its Cl^- salt. The view is the same as in Figure 1c but the oxygen and nitrogen atoms are now represented by medium-sized shaded spheres, and sulfur, carbon and hydrogen atoms are represented by large, medium and small open spheres, respectively.
- Figure 1e. shows a space-filling drawing of the structure for the $[C_{27}H_{36}N_3O_3S]^+$ cation, as observed in the solid-state structure of its Cl^- salt. The view is as in Figure 1c.

Exhibit Z

Figure 1f. shows a perspective drawing of the hydrogen-bonding interaction involving the Cl^- anion and the amine (N_1 and N_2) nitrogens of two symmetry-related $[\text{C}_{27}\text{H}_{36}\text{N}_3\text{O}_3\text{S}]^+$ cations in the crystal. Atoms of the cation related by symmetry operation $-x, 1-y, -0.5+z$ to the one shown in Figure 1d are labeled with a prime ('). Atoms are represented as in Figure 1b and the hydrogen bonds between the Cl^- anion and the $[\text{C}_{27}\text{H}_{36}\text{N}_3\text{O}_3\text{S}]^+$ cations are shown with dashed lines.

Figure 1g. shows a perspective drawing of the hydrogen-bonding interaction involving sulfonamide proton $\text{H}_{3\text{N}}$ and the sulfonamide oxygen atom of a symmetry-related $[\text{C}_{27}\text{H}_{36}\text{N}_3\text{O}_3\text{S}]^+$ cation in crystals of $[\text{C}_{27}\text{H}_{36}\text{N}_3\text{O}_3\text{S}][\text{Cl}]$. Atoms of the $[\text{C}_{27}\text{H}_{36}\text{N}_3\text{O}_3\text{S}]^+$ cation related by symmetry operation $x, 1+y, z$ to the one shown in Figure 1d are labeled with double primes ("). Atoms are represented as in Figure 1b and the hydrogen bond between $[\text{C}_{27}\text{H}_{36}\text{N}_3\text{O}_3\text{S}]^+$ cations is shown with a dashed line as are the hydrogen bonds between the cation and Cl^- anion within each of the two asymmetric units shown.

Exhibit AB

cyclicAmidines



























	JNJ 6141694 BID 3498187 salt AAC NB 14794 page 5 # 1		JNJ 6141694 BID 4251723 salt AAC NB 8902 page 949 # 1
	JNJ 6141694 BID 5855332 salt AAA NB 17472 page 2 # BHF		JNJ 7494513 BID 4323508 salt AAA NB 14794 page 96 # 1
	JNJ 7572045 BID 4362274 salt AAA NB 14794 page 145 # 1		JNJ 7572071 BID 4362288 salt AFP NB 14794 page 147 # 1
	JNJ 8283535 BID 4730292 salt AAC NB 14794 page 130 # 1		JNJ 8283626 BID 4730334 salt AAC NB 14794 page 131 # 1
	JNJ 8487960 BID 4832499 salt AAC NB 14811 page 138 # 0		JNJ 8512335 BID 4844686 salt AAC NB 14794 page 135 # 1
	JNJ 8702460 BID 5213677 salt AAA NB 17472 page 2 # BGW		JNJ 8702460 BID 9713956 salt AAC NB 8902 page 946 # 1
	JNJ 17225728 BID 9890034 salt AAA NB 14794 page 57 # 1		JNJ 17225767 BID 9890048 salt AAA NB 14794 page 62 # 1

Exhibit AC

	JNJ 17225858 BID 9890097 salt AAC NB 14794 page 65 # 1		JNJ 17225858 BID 9943472 salt AAC NB 14794 page 87 # 1
	JNJ 17225858 BID 9943717 salt AAC NB 14794 page 150 # 1		JNJ 17225884 BID 9890104 salt AAC NB 14794 page 70 # 1
	JNJ 17228107 BID 9891224 salt AAA NB 14794 page 75 # 2		JNJ 17228276 BID 9891308 salt AAC NB 14794 page 84 # 1
	JNJ 17231019 BID 9892680 salt AFP NB 14794 page 143 # 1		JNJ 17231318 BID 9892827 salt AFP NB 14794 page 151 # 1
	JNJ 17825522 BID 10358124 salt AAC NB 14794 page 99 # 1		JNJ 17825548 BID 10358131 salt AAC NB 14794 page 100 # 1
	JNJ 17988906 BID 10448627 salt AFP NB 17182 page 124 # 2		JNJ 17988919 BID 10448634 salt AFP NB 17182 page 124 # 1

Project No. *NPY5*Protocol / Experiment No. *308*



Date

Subject


Purpose

Contd. from page

REAL NUMBER	Properties	NPY5 IC50 nM
277824-300-A	Y5	38
277825-300-A	Y5	3000
277826-300-A	Y5	1100
277827-300-A	Y5	621
277833-300-A	Y5	29
277802-300-A	Y5	422
<i>1277512-300-A</i>	Y5	417
<i>279554-300-A</i>	Y5	437
057926-300-C	OB/OB	
058532-300-A	OB/OB	
058746-300-A	Y2,Y1	
058747-300-A	Y2	
058748-300-A	Y2	
058749-300-A	Y2	
275087-300-A	Y2	
277829-300-A	Y2	
58260-300-A	CONTROL	9
58442-300-A	CONTROL	31
58660-300-A	REPEAT	661
58691-300-A	REPEAT	225

Binding of new RWJ
compounds to recombinant
human NPY5 receptorBinding of new RWJ
compounds to recombinant
human NPY5 receptor

Structure	Properties	Samples
RWJ Number 277833-300-A		
Entry Date		
Site Springhouse		
Team CNS Research		
Prep. by/Approval Nicole Willard		
Notebook/Page/Id 14779-62-1		
Supplier PRI SPRING HOUSE		
MW 482.6852	Batch MW 519.148	CAS Cat#
Formula C ₂₈ H ₃₈ N ₂ O ₃ S		Batch Formula C ₂₈ H ₃₈ N ₂ O ₃ S.HCl
Name		



RWJ Number: 277824-300-A

Mol. Form: C₂₈H₃₈N₂O₃S

Mol. Wt.: 451.63

*Could be
very
interesting**Jim must confirm
structure*

Contd. on page

Investigator *[Signature]*

Date

Read and Understood *[Signature]*


Date

Project No. NPY5 Protocol / Experiment No. 311 Date

Subject Purpose


Contd. from page

Binding of new RWJ
compounds to recombinant
human NPY5 receptor



REAL NUMBER	NPY5 IC50 nM
289060-300-A	324
289062-300-A	33
289136-300-A	40
289137-300-A	123
279518-300-A	37
279527-300-A	38
058532-300-A	
279527-300-A	
277824-300-B	26
277824-300-A	40
058260-300-A	6
058442-300-A	23
058669-300-A	17

Binding of new RWJ
compounds to recombinant
human NPY5 receptor



Contd. on page

Investigator [Signature] Date


Read and Understood [Signature] Date

Project No. NPY5 Protocol / Experiment No. 320 Date:

Subject Purpose

Contd. from page

Data 1	Data2	Data3	Average	Specific	%Specific	%Inhibition	REAL NUMBER	NPY5 IC50 nM
1772.6	1866.3	297521	1819	19	0	100	297521-300-A	62
2072.1	2159.4		2116	316	5	95	297561-300-A	106
2852.5	3285.6		3069	1269	20	80	297633-300-A	2
3838.7	3889.2		3864	2064	33	67	297679-300-A	150
5800	6117.2		5959	4159	67	33	295986-300-A	39
7512.8	7430.1		7471	5671	91	9	293708-300-A	Inactive
7163	7681.6		7422	5622	91	9	293710-300-A	Inactive
7749.2	8067.8		7909	6109	99	1	296148-300-A	
1890.4	1784.1	297561	1837	37	1	99	058427-300-A	29
2218.7	2058.1		2138	338	45	95	058427-300-C	35
2844.8	2981.9		2913	1113	18	82	293376-300-A	39
4829.7	4741.8		4786	2986	48	52	293407-300-A	73
6403.8	6567.5		6486	4686	76	24	293420-300-A	362
7598.2	7494.5		7547	5747	93	7	293377-300-A	50
7484.8	7733.1		7614	5814	94	6	303114-300-A	412
7619.7	7641.2		7580	5780	93	7	303115-300-A	15
2017.8	1777.3	297633	1897	97	2	98	058442-300-A	29
2262.4	2151.4		2207	407	7	93	058260-300-A	8
2140.9	2121.3		2131	331	5	95	WRY-amide	3400
2161.2	2249		2205	405	7	93		
2538.5	2250.7		2395	595	10	90		
2946.8	2914		2930	1130	18	82		
4846.8	4301.6		4574	2774	45	55		
6667.8	6814.3		6741	4941	80	20		
2047.1	1892.8	297679	1970	170	3	97		
2238.8	2306.5		2273	473	8	92		
3152.7	3632.1		3392	1592	28	74		
5382.4	5383.6		5383	3583	58	42		
6760.8	7085.7		6923	5123	83	17		
8191.5	7565		7878	6078	98	2		
7928.7	8018.8		7974	6174	100	0		
7400.3	7230.7		7318	5516	89	11		
1752.5	1832.5	58260	1793	-8	0	100		
2082.8	2221.3		2152	352	6	94		
2100.7	1872.1		1986	186	3	97		
2215.8	2305.3		2261	461	7	93		
2733.8	3076.5		2905	1105	18	82		
4252.1	4826.2		4589	2789	45	55		
6222.1	6488.7		6345	4545	73	27		
7283.4	6737.6		7011	5211	84	16		
6418.8	5552.1	WRY-amide	5486	3686	59	41		
6808.8	8742.3		6775	4975	80	20		
7898	7328.1		7613	5813	94	6		
7886.1	7543.1		7720	5920	95	5		
8281.8	7680.8		7986	6188	100	0		
8337.7	7927.3		8133	6333	102	-2		
8154.8	9083.6		8619	6819	110	-10		
7288.4	7134.5	Control	7201	5401	87	13		



RW 297633

Came out at less than 2m.

14794-65-1
Molecular Weight = 518.12

Contd. on page

Investigator [Signature]

Date



Read and Understood [Signature]

Date

Project No. NPY5 Protocol / Experiment No. 323 Date _____

Subject _____ Purpose _____

Contd. from page _____

Binding of new RWJ
compounds to recombinant
human NPY5 receptorBinding of new RWJ
compounds to recombinant
human NPY5 receptor

Compound	REAL NUMBER	NPY5 IC50 nM
14811-55	298270-300-A	Y2
14811-57	303090-300-A	Inactive
14811-72		227
14811-74	306450-300-A	Y2
14794-75-2	303088-300-A	44 ✓
14794-76-1	058260-300-C	8
14794-84-1	303193-300-A	58
14794-87-1	297633-300-B	47 ✓
	058260-300-C	Large Scale
	297633-300-B	Large Scale ✓
REPEATS/ CONTROLS	297521-300-A	58 ✓
	297679-300-A	125 ✓
	297561-300-A	76 ✓
	058260-300-A	4
	058442-300-A	16
	297633-300-A	1.5 ✓

It turns out I made mistake by
switching caps.

297633 was switched w/ 297679 so the

297633-300-B data is accurate

297633-300-A is actually 125 nM

297679-300-A is actually 1.5 nM

Contd. on page _____

Investigator _____

Date _____


Read and Understood _____

Date _____


Subject **Purpose**

Contd. from page _____

NUMBER	Properties	NPY5	NPY2
		IC50 nM	IC50 nM
66-300-A	Y2		Inactive
13-300-A	Y2		Inactive**
99-300-A	Y2		3000
00-300-A	Y2		10000
22-300-A	Y5	133	
52-300-A	Y5	Inactive	
51-300-A	Y5	Inactive	
39-300-A	Y5	111	
40-300-A	Y5	54	
41-300-A	Y5	9	
43-300-A	Y5	647	
44-300-A	Y5	1000	
55-300-A	Y5	200	
60-300-A		3	
79-300-A		1	
88-300-A		67	
02-301-A			3000
71-301-A			3000
43-300-A	This is the cyclopropyl replaceme		4000





13 showed no dose response
overall lower amount of binding




Contd. on page _____ Investigator [Signature] Date _____


Read and Understood S. K. K. K. Date

Project No. NPY5 Protocol / Experiment No. 342 Date Subject Purpose Contd. from page 171Binding of new RWJ
compounds to recombinant
human NPY5 receptorBinding of new RWJ
compounds to recombinant
human NPY5 receptor

Compound	REAL NUMBER	NPY5 IC50 nM
14794-111-1	313746-300-A	23
14794-112-1	313747-300-A	2000
14794-113-1	313748-300-A	20
14205-087A	313682-300-A	769
14205-087-B	313683-300-A	133
14205-086-A	313684-300-A	745
14205-086-B	313685-300-A	200
14811-101	313664-300-A	200
14811-103	313665-300-A	7
14811-106	313725-300-A	33
14811-107-1	313717-300-A	142
Control	297679-300-A	1

14811-103
RWJ 313665-300-A

Here is another one:

14811-106
RWJ 313725-300-A


Removal of cyclopropyl salt
did not hurt activity when replaced
with amide

Contd. on page Investigator Date Read and Understood Date

Project No. NPY5 Protocol / Experiment No. 348 Date
 Subject Purpose

Contd. from page


Binding of new RWJ compounds to recombinant human NPY5 receptor




314465
 314466
 314467
 314464
 297679
 58442

	IC50
313725	96
313717	286
314246	175
314247	2500
314463	807
58260	6
314465	75
314466	23
314467	54
314464	548
297679	1
58442	22

Binding of new RWJ compounds to recombinant human NPY5 receptor



313725
 313717
 314246
 314247
 314463
 58260



Contd. on page

Investigator

Date

Read and Understood

Date


No. 14280-187

Project No. NKPS Protocol / Experiment No. 351 Date _____


Subject _____ Purpose _____

Contd. from page _____

	Data2	Data3		Average	Specific	%Specific	%Inhibition
1730.6	1890.9		313665	1711	-10	0	100
1728.8	1796.3			1763	41	1	99
2000.5	2497.4			2249	528	9	91
2742.8	2744.2			2744	1022	18	82
3990.1	3850.5			3920	2199	39	61
5146.5	5760.6			5454	3732	66	34
6762.2	6308.8			6536	4814	86	14
7505	7525.8			7515	5794	103	-3
1882.3	1913.9		314246	1898	177	3	97
2035.5	2313.7			2175	453	8	92
3535.9	3335.2			3436	1714	31	69
5712.5	5090.1			5401	3680	66	34
6807.5	7315.8			7062	5340	95	5
7195.1	7967.3			7581	5860	104	-4
7219.6	7832.4			7526	5805	103	-3
7844.8	8201.4			8023	6302	112	-12
3708.3	3404.6		314247	3555	1834	33	67
5258.8	5099.2			5178	3457	62	38
6904	6586.6			6745	5024	89	11
6042.5	8084.2			7063	5342	95	5
7762.7	7409.5			7586	5865	104	-4
8388.4	7608.5			7987	6266	112	-12
7357.8	7202.5			7285	5564	99	1
7768.7	8088.2			7928	6207	111	-11
1682	1953.5		313725	1818	97	2	98
1883.9	1989.6			1987	265	5	95
2491.4	2742.3			2617	896	16	84
3935.4	3815.9			3876	2154	38	62
6172.9	5789.1			5981	4260	76	24
7587.8	7311.9			7440	5719	102	-2
7445.1	7746.6			7596	5875	105	-5
8394.5				8395	6673	119	-19
2013.3	1917.6		313717	1965	244	4	96
2672.5	3248.2			2960	1239	22	78
4246.7	4076.5			4162	2440	43	57
5751.1	5745.9			5749	4027	72	28
6869.2	7102.1			6986	5264	94	6
7480.5	7683.5			7582	5861	104	-4
7240.7	8086.7			7664	5943	106	-6
7941.2	7990.5			7966	6245	111	-11
1871.6	1770.8		58260	1721	0	0	100
1886	1681.1			1784	62	1	99
2076.7	2341.8			2209	488	9	91
2880.3	2144.6			2517	796	14	86
2884.4	2708.1			2786	1075	19	81
3455.5	3402.1			3429	1708	30	70
5312.5	4871.2			5142	3421	61	39
7279	7394.3		Control	7337	5615	100	0



Data 1	Data2	Data3		Average	Specific	%Specific	%Inhibition
2456.6	2430.7		314704	2444	544	11	89
2739.4	2900.1			2820	920	18	82
4278.8	4366.8			4323	2423	48	52
5951.1	6528.8			6240	4340	85	15
6717.5	7331.8			7025	5125	100	0
6924.7	7385.2			7155	5255	103	-3
6572.3	6949.6			6781	4861	95	5
7262.5	7476.5			7370	5470	107	-7
1937.2	1741.6		316980	1839	-61	-1	101
2685.5	2620.7			2653	753	15	85
3695.3	3290.6			3493	1593	31	69
5036.2	5036.8			5037	3137	62	39
6449.1	6933.5			6541	4841	91	9
7041.7	7184.3			7113	5213	102	-2
	7594.3			7594	5694	112	-12
7424.7	7584.5			7505	5605	110	-10
4451.5	4468.6		317050	4460	2560	50	60
5900.7	6485.8			6193	4293	84	16
7050.1	7051.4			7051	5151	101	-1
7207.2	8055.1			7631	5731	112	-12
7538.5	7933.9			7736	5836	114	-14
7205.9	7553.6			7380	5480	107	-7
6998.1	7893.8			7448	5546	109	-9
7424.8	7996.7			7711	5811	114	-14
2954.4	3050		317051	3002	1102	22	78
4380	4226.5			4303	2403	47	53
6070.8	5922			5996	4096	80	20
8136.1	7411			7774	5874	115	-15
8036.8	7825.4			7932	6032	118	-18
7312.3	7286.3			7299	5399	106	-6
7481.5	7281.5			7382	5482	107	-7
7261.6	7254.6			7258	5358	105	-5
1797.9	1996.7		314503	1897	-3	0	100
1959.9	2263			2111	211	4	98
2461.5	2471			2468	566	11	89
3870.7	3731			3801	1901	37	63
5563.8	5263.4			5414	3514	69	31
6611.4	6575.5			6593	4693	92	8
6761.6	6793			6777	4877	96	4
7257	7072.9			7165	5265	103	-3
1948.2	1781.9		314709	1865	-35	-1	101
2329.3	2353.7			2342	442	9	91
2701.3	2585.5			2643	743	15	85
3866.2	3959.8			3913	2013	39	61
5614	5543.3			5579	3679	72	28
7131.5	6330.6			6731	4831	95	5
7147.7	8214.3			7681	5781	113	-13
7019.4	7358.9		Control	7189	5289	104	-4



Contd. on page _____



Investigator _____

Date _____

Project No. NPY5 Protocol / Experiment No. 357 Date

Subject Purpose

Contd. from page

Binding of new RWJ
compounds to recombinant
human NPY5 receptorBinding of new RWJ
compounds to recombinant
human NPY5 receptor

REAL NUMBER	NPY5 IC50 nM
320563-300-A	32
320564-300-A	8
320566-300-A	Inactive
320605-300-A	878
318251-300-A	117
318252-300-A	539
320112-300-A	202
057836-300-A	222
318245-300-A	530
316035-300-A	168
314996-300-A	8
297679-300-A	1.7
314995-300-A	1

Contd. on page Investigator Date

Read and Understood S. Kuan Date

Project No. NPY5D Protocol / Experiment No. 369 Date


Subject Purpose

Contd. from page

Data 1	Data2	Data3	Average	Specific	%Specific	%Inhibition
7271.3	7423.3	331781	7347	5737	131	-31
5374.9	5799.9		5587	3977	91	9
5953.6	5857.9		5906	4296	98	2
5878.1	6608.2		6243	4633	106	-6
5701.8	5824.3		5763	4153	95	5
6088.1	6384		6236	4626	105	-5
6240	6356.7		6298	4688	107	-7
6526.3	6016.1		6271	4661	106	-6
3694.8	3622.7	332260	3659	2049	47	53
2998	4601.5		3800	2190	50	50
5252.6	6001		5627	4017	91	9
5613	5870.8		5742	4132	94	6
5945.7	5986.8		5966	4356	99	-1
6164.7	6373.3		6269	4659	106	-6
6588.9	6653.7		6611	5001	114	-14
6316.8	6139.6		6228	4618	105	-5
1626.5	1590.9	332517	1609	-1	0	100
1768.1	1665.1		1717	107	2	98
2145.3	1894.5		2020	410	9	91
2659.2	2555.6		2607	997	23	77
4092.4	4004.6		4049	2439	56	44
5542.6	5123.9		5333	3723	85	15
5744.3	5893		5819	4209	96	4
6199.3	5989.4		6094	4484	102	-2
2067.4	2437.3	332259	2252	642	15	85
2395.7	2665.3		2531	921	21	79
3591.6	4169.9		3881	2271	52	48
5104.9	5220.7		5163	3553	81	19
5744.7	6313.6		6029	4419	101	-1
6673.9	5753		6213	4603	105	-5
6504.6	6073.6		6289	4679	107	-7
6315.8	6306.9		6311	4701	107	-7
4490.8	4745.9	331774	4618	3008	69	31
5330.2	5577.4		5454	3844	88	12
5932.1	6394.9		6164	4554	104	-4
5985	6198.6		6092	4482	102	-2
6143.8	6434.1		6289	4679	107	-7
6132.9	5790.3		5962	4352	99	1
6173.1	5604.6		5889	4279	97	3
5865.8	5541.5		5704	4094	93	7
1756.9	1514.9	332261	1636	26	1	99
2869.7	1848		2359	749	17	83
2341	2601.6		2471	861	20	80
4248.5	3953.2		4101	2491	57	43
5654.1	5470.7		5562	3952	90	10
5457.5	6063.7		5761	4151	95	5
6348.8	6425.1		6387	4777	109	-9
5996.3	5644.6		5820	4210	96	4

REAL NUMBER	NPY5 IC50 nM
332261-300-A	121 ✓
332259-300-A	263 ✓
332260-300-A	3000
331781-300-A	>3000
331774-300-A	>3000
331775-300-A	54
332285-300-A	>3000
332360-300-A	2000
332300-300-A	>3000
114102-301-B	Y2
332517-300-A	35 ✓
332516-300-A	Y2
332518-300-A	2000
328568-300-A	265
306468-300-A	250
313717-300-A	280
328577-300-A	856
313665-300-A	39
314613-300-A	240
306672-300-A	133

Numbers are a bit higher than normal.

↓
Repeat

Contd. on page

Investigator

Date

Read and Understood

Date

Project No. NPY5Protocol / Experiment No. 374


Date

Subject

Purpose


Contd. from page

Binding of RWJ compounds to
recombinant human NPY5
receptor.



Binding affinities
were similar for
both rat +
human receptor

Binding of RWJ compounds to
recombinant rat NPY5 receptor.



Contd. on page

Investigator

Date



Read and Understood

Date

Project No. NPY5 Protocol / Experiment No. 388 Date

Subject Purpose

Contd. from page

Binding of new RWJ
compounds to recombinant
human NPY5 receptorBinding of new RWJ
compounds to recombinant
human NPY5 receptor

RWJ NUMBER	NPY5 IC50 nM
334170-300-A	335
333329-300-A	to be repeated
333328-300-A	Inactive
333592-300-A	2000
333319-300-A	Inactive
333591-300-A	2000
333590-300-A	836
058669-000-B	7
332556-300-A	1000
058442-002-A	
058669-300-D	
058442-300-A	11
297679-300-A	1.5
058260-300-A	4

Contd. on page

Investigator

Date

Read and Understood

Date

No. 15229- 46

Project No. NPYProtocol / Experiment No. 385

Date

Subject

Purpose

Contd. from page

Data 1	Data2	Data3	Average	Specific	%Specific	%Inhibition	Data 1	Data2	Data3	Average	Specific	%Specific	%Inhibition
983.4	861.7		296679	923	88	3	1687.6	1772		333592	1730	844	37
849.6	823.4			837	2	0	2461.3	2536.1			2499	1713	68
975.9	1084.2			1030	195	7	3135.9	3279.3			3208	2422	96
1046.6	1018.4			1033	198	7	3478.2	3778			3628	2843	112
1093.4	1092.4			1093	258	10	3542.1	3337			3440	2654	105
1161.6	1386.7			1274	439	17	3536.8	3692.9			3615	2829	112
2019.2	1991			2005	1170	44	3547.2	3489.4			3518	2733	108
2803.4	2994.6			2899	2064	78	3508.5	3706.7			3608	2822	111
934.9	989.6		58442	962	127	5	6302.6	2225.2		333319	4264	3478	137
975.4	748.3			862	27	1	3316.3	3142.6			3229	2444	96
1057.4	1010			1034	199	8	3549.3	3500.9			3525	2740	108
1261.5	1040.1			1151	316	12	3626.9	3714			3670	2885	114
1649.2	1616			1633	798	30	3552	3873			3713	2927	115
2523.9	2586.7			2555	1720	65	3848.8	3985.8			3917	3132	123
3257	3123.4			3190	2355	89	3610.6	3514.8			3563	2777	110
3856.8	3380.8			3619	2784	106	3596.4	3561.1			3579	2793	110
885.2	802.5		334170	844	9	0	1555.4	1466.1		333591	1511	725	29
1136.5	1138.3			1137	302	11	2316.5	2663.8			2490	1705	67
2189.1	2074.5			2132	1297	49	3533.4	3340.9			3437	2652	105
3087.3	3050.6			3069	2234	85	3603.7	3471.4			3538	2752	109
3532.9	3558.4			3546	2711	103	3694.4	3303.1			3499	2713	107
3764	3539.1			3652	2817	107	3627.2	3700.2			3664	2878	113
3774.1	3618.7			3696	2861	109	3704.3	3673.1			3689	2803	114
3612.3	3697.4			3655	2820	107	3710.7	3923.9			3817	3032	120
2867.9	3213.8		333329	3041	2206	84	1060.6	1164.6		333590	1113	327	13
3777.6	800			2289	1454	55	1671.4	1626.9			1649	864	34
4170.6	1203.1			2687	1852	70	3200	2702.5			2951	2166	85
3943.3	1546.7			2745	1910	72	3148.2	3157.4			3153	2367	93
3740.9	2130.6			2936	2101	80	3542.3	3363.7			3453	2668	105
3946.5	2992.3			3469	2634	100	3609.6	3645.6			3628	2842	112
3792.8	3364.9			3579	2744	104	3676.6	3482.5			3581	2795	110
3865.1	3664.8			3765	2930	111	3713.6	3345			3529	2744	108
3440.6	3190.3		333328	3315	2480	94	1549.3	1529.5		332556	1539	754	30
3485.2	3792.4			3639	2804	106	2407.1	2157.8			2282	1497	59
3760.8	3731.9			3746	2911	110	3066.6	2977.8			3022	2237	88
3602.8	4130.1			3866	3031	115	3468.6	3079.8			3274	2489	98
4474.9	3668.3			4072	3237	123	3374.2	3251.9			3313	2528	100
3920	3725.3			3823	2989	113	3470	3006.6			3238	2453	97
3677.5	3601.8			3640	2805	106	3578.8	3465.6			3522	2737	108
3648.8	3709.3			3679	2844	108	3505.9	3806			3656	2870	113
792.9	877.1		58260	835	0	0	771.8	799.2		58669	786	0	0
866	770.3			818	-17	-1	1012	997.2			1005	219	9
998.2	978.9			989	154	6	968.5	904.7			937	151	6
1080.3	1085.7			1083	248	9	1209.9	1085.2			1148	362	14
1099.8	1267.6			1184	349	13	1270	1371.6			1321	535	21
1549	1758.8			1654	819	31	2044.8	1874.1			1959	1174	46
2612.3	3033.5			2823	1988	75	3000	2867.6			2934	2148	85
3120.4	3823.6		Control	3472	2637	100	3293.4	3349.6		Control	3322	2536	100

Contd. on page

Investigator

Date



Read and Understood

Date

Project No. NPY5 Protocol / Experiment No. 465 Date _____

Subject _____ Purpose _____

Contd. from page _____

Binding of new compounds to
NPY5 receptorBinding of new compounds to
NPY5 receptorContd. on page _____ Investigator [Signature] Date _____Read and Understood [Signature] Date 7